

## SHORT NOTES

## Non-breeding reed warblers *Acrocephalus scirpaceus* (Hermann, 1804) in June in southern Europe: local or still migrating birds?

Juan Arizaga<sup>1,2\*</sup>, Agustín Mendiburu<sup>2</sup> & Daniel Alonso<sup>2</sup>

<sup>1</sup> Institute of Avian Research "Vogelwarte Helgoland", an der Vogelwarte 21, 26386 Wilhelmshaven, Germany.

<sup>2</sup> Aranzadi Ringing Scheme, Aranzadi Sciences Society, Zorroagaina 11, 20014 San Sebastián, Spain.

\* Corresponding author: juan.arizaga@ifv-vogelwarte.de

KEY WORDS: Bird breeding season, floaters, flight morphology, population structure, Txingudi.

Accurate estimations of population structures for breeding birds are essential for assessing population-specific parameters, such as adult survival rates, breeding site fidelity or productivity (1-3). However, not all adult birds observed in a certain breeding area should be considered as breeders: non-breeding local birds and, in migrant species, individuals that are still migrating could also be present. Taking the proportions of birds in these categories into account is crucial for estimating unbiased population-specific parameters (4; 5).

In migrant bird species with large distribution areas, the overlap between local breeding birds and still-migrating ones is frequent, particularly at low latitudes (6-8). Compared to non-migrants (e.g., breeding birds), migrants normally show higher mean fuel loadings, i.e., higher energy stores that are used during the migratory flights (9). Birds tend to have longer and more pointed wings as well as shorter tails related to body size with increasing migration distances (10; 11) as is confirmed in several species of Holarctic passerines (12-15). Therefore, both a higher mean fuel load and different morphologies more resembling the long-distance-type in non-breeders may indicate that these birds are not local but still migrating birds.

Floaters present the non-territorial fraction of a territorial population (16) and are often older, less-experienced or handicapped birds (17), that would normally not breed. Because of their transient behaviour, they are less likely to be recaptured (4). The reed warbler *Acrocephalus scirpaceus* (Hermann, 1804) is a widespread breeding songbird in western Europe that passes the winter in tropical Africa (7). In spring, reed warblers reach southern Europe by late-March or April (7). Thus, most (if not all) adults should have reached their breeding areas by June, at least in southern Europe (18). The question whether some northern European birds that are still migrating could be found in southern European breeding areas has rarely been investigated. Our aim here was to determine whether non-breeding reed warblers captured in June in southern Europe were local or migrating birds, by considering traits such as fuel load, flight-associated morphology and recaptures.

Reed warblers were captured in Jaizubia (43°21'N, 01°49'W, Fuenterrabía, N Iberia), a 38.1ha small tidal

river area located in the Txingudi marshlands. Birds were captured with mist nets (240 linear m) placed across reed beds and opened for four hours starting at dawn, during 10 consecutive days in June (05.06–14.06), July (10.07–19.07), and August (07.08–16.08) 2006. Once captured, reed warblers were ringed (or the ring was read) and their age was determined as adults (second-year birds or older, EURING code 4) or juveniles (first-year birds, EURING code 3). The following variables were also recorded: wing length (method III,  $\pm 0.5$ mm; 19), tail length ( $\pm 0.5$ mm), length of primary feathers P2 to P9 (from outermost to innermost,  $\pm 0.5$ mm; P1 is vestigial), body mass (TANITA digital balance,  $\pm 0.1$ g) and fat score (scaled from 0 to 8; 20). Finally, we checked whether birds had brood patches (females; the brood patch is used for incubating the eggs, has no feathers and is highly vascularized) or cloacal protuberances (males, scaled from 0 to 9 (21)). We only investigated adult birds because our aim was to analyse the characteristics of non-breeding birds in a breeding population.

Flight-associated morphology (wing length, wing morphology, tail length/wing length) and fuel load (body mass/wing length and fat score; here, wing length is used as a proxy of body size; 22) were compared between breeding and non-breeding adult reed warblers captured in June. A bird was considered as breeder if it showed apparent breeding signs in June, July, or August. Because data on flight-associated morphology and body mass/wing length fitted the normal distribution (K-S test,  $P > 0.05$ ),  $t$ -tests were used for statistical comparisons. In contrast, fat score was not normally distributed (K-S test:  $Z = 2.466$ ,  $P < 0.001$ ), which is why a  $U$  test was applied. In order to avoid pseudo-replication, each individual was considered only once.

Wing morphology was assessed by the  $C_2$  index which expresses wing pointedness (23). Positive values of the index are associated with rounded wings, while negative values are associated with more pointed wings.

Of the 77 adults captured in June, 47 (61.0%) showed signs of breeding (either in June or later, in July or August), and were considered to be breeders. This value is similar to that found in some Nearctic passerines (4) and reinforces the idea that a significant fraction of adult birds in a certain local population consists of non-breeders during the breeding season (24).

Breeding and non-breeding reed warblers captured in June had the same flight-associated morphology but differed in relation to their fuel load (Table 1). This supports

the hypothesis that non-breeders may be either local birds or non-local birds from populations breeding in areas relatively close to Txingudi, because if these were birds from more northern areas, we would expect more pronounced long-distance morphology (13). A higher fuel load in non-breeders may indicate that they are still migrating birds breeding in areas close to Txingudi. Alternatively, these individuals could be local, either birds arriving later in their breeding areas (with some remaining fuel accumulated for the spring migration period, from Africa to Europe) or birds starting to gain the fuel required for the migration beginning in summer (from Europe to Africa).

None of the non-breeding birds captured in June was recaptured afterwards, whereas 40.4% of breeders at Txingudi were recaptured in July or August ( $\chi^2_1=16.101$ ,  $P$ -exact<0.001; Table 2). This confirms that the recapture probability was relatively high throughout the season and suggests that non-breeders did not remain at Txingudi for a long time. This result is consistent with the idea that non-breeders could either still be migration birds or local transient birds (floaters), unable to defend a territory (25; 4).

The presence of non-breeders can be very important at a local level, and in our opinion, this should be considered in studies dealing with the dynamics of breeding bird populations. This is particularly relevant in programs surveying parameters such as annual survival, population size or productivity (1). All of these parameters are fundamental to our understanding of population trends in particular species and, therefore, are also directly linked to the conservation of biodiversity.

TABLE 1

Differences in flight-associated biometrics and fuel load (mean±SE) between breeding and non-breeding adult reed warblers captured in June in the Txingudi marshlands (northern Iberia).

| Variable                    | Breeding            | Non-breeding        | Statistics     | P      |
|-----------------------------|---------------------|---------------------|----------------|--------|
| Wing length                 | 63.9±0.3<br>(N=47)  | 63.9±0.4<br>(N=30)  | $t_{75}=0.011$ | 0.991  |
| Wing morphology (roundness) | 0.75±0.02<br>(N=43) | 0.73±0.02<br>(N=29) | $t_{70}=0.948$ | 0.346  |
| Tail length/Wing length     | 0.84±0.01<br>(N=47) | 0.84±0.01<br>(N=30) | $t_{75}=0.428$ | 0.670  |
| Body mass/Wing length       | 0.16±0.01<br>(N=47) | 0.17±0.01<br>(N=30) | $t_{75}=2.830$ | 0.006  |
| Fat scores                  | 0.7±0.1<br>(N=43)   | 2.5±0.2<br>(N=29)   | $U=199.0$      | <0.001 |

TABLE 2

Number of breeding and non-breeding adult reed warblers captured at Txingudi in June and recaptured at the same site in July or August.

|              | Captured only in June | Captured in June and July/August |
|--------------|-----------------------|----------------------------------|
| Breeding     | 28                    | 19                               |
| Non-breeding | 30                    | 0                                |

## ACKNOWLEDGEMENTS

Thanks to all the people who collaborated during the field work, especially to J. I. Jauregi, R. Martín and J. M. Sánchez. The Diputación de Gipuzkoa authorized us to conduct the ringing activities. J. A. was supported by a post-graduate fellowship from the Gobierno Vasco – Eusko Jaurlaritz. The Gobierno Vasco – Eusko Jaurlaritz founded the project. V. Salewski, G. R. Geupel and two anonymous referees provided very valuable comments that improved an earlier version of this work.

## REFERENCES

1. PEACH WJ, BUCKLAND ST & BAILLIE SR (1996). The use of constant effort mist-netting to measure between-year changes in the abundance and productivity of common passerines. *Bird Study*, 43:142-156.
2. PEACH WJ, BAILLIE SR & BALMER DE (1998). Long-term changes in the abundance of passerines in Britain and Ireland as measured by constant effort mist-netting. *Bird Study*, 45:257-275.
3. MILES W, FREEMAN SN, HARRISON NM & BALMER DE (2007). Measuring passerine productivity using constant effort sites: the effect of missed visits. *Ring and Migration*, 23:231-237.
4. NUR N, GEUPEL GR & BALLARD G (2000). The use of constant-effort mist-netting to monitor demographic processes in passerine birds: annual variation in survival, productivity and floaters. In: BONNEY R, PASHLEY DN, COOPER RJ & NILES L (eds), 3rd Partners in Flight workshop, U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Center, Ogden, UT.: 185-194.
5. SÆTHER BE, LANDE R, ENGEN S, WEIMERSKIRCH H, LILLEGAAARD M, ALTWEGG R, BECKER PH, BREGNBALLE T, BROMMER JE, MCCLEERY R, MERILA J, NYHOLM E, RENDELL W, ROBERTSON RR, TRYJANOWSKI P & VISSER ME (2005). Generation time and temporal scaling of bird population dynamics. *Nature*, 436:99-102.
6. CRAMP S (1988). *Handbook of the Birds of Europe, the Middle East and North Africa*. Vol. 5. Oxford University Press, Oxford.
7. CRAMP S (1992). *Handbook of the Birds of Europe, the Middle East and North Africa*. Vol. 6. Oxford University Press, Oxford.
8. CRAMP S & PERRINS CM (1994). *Handbook of the Birds of Europe, the Middle East and North Africa*. Vol. 9. Oxford University Press, Oxford.
9. NEWTON I (2008). *The migration ecology of birds*. Academic Press, London.
10. MÖNKKÖNEN M (1995). Do migrant birds have more pointed wings?: a comparative study. *Evolutionary Ecology*, 9:520-528.
11. CALMAESTRA R & MORENO E (2001). A phylogenetically-based analysis on the relationship between wing morphology and migratory behaviour in passeriformes. *Ardea*, 89:407-416.
12. CHANDLER CR & MULVIHILL RS (1990). Interpreting differential timing of capture of sex classes during spring migration. *Journal of Field Ornithology*, 61:85-89.
13. GARCÍA-PEIRÓ I (2003). Intraspecific variation in the wing shape of the long-distance migrant Reed Warbler *Acrocephalus scirpaceus*: effects of age and distance of migration. *Ardeola*, 50:31-37.

14. CHERNETSOV N (2004). Intraspecific variation of wing pointness index in juvenile *Acrocephalus* warblers in the south-eastern Baltic. *Journal of Ornithology*, 145:105-108.
15. FIEDLER W (2005). Ecomorphology of the external flight apparatus of Blackcaps *Sylvia atricapilla* with different migration behavior. *Annals of the New York Academy of Sciences*, 1046:253-263.
16. WINKER K (1998). The concept of floater. *Ornitología Neotropical*, 9.
17. PODULKA S, ROHRBAUGH RW & BONNEY R (2004). *Handbook of bird biology*. Cornell Lab. of Ornithology, New Jersey.
18. BERMEJO A (2004). *Migratología, estructura y dinámica poblaciones de paseriformes asociados a vegetación de ribera*. Thesis. Universidad Autónoma de Madrid, Madrid.
19. SVENSSON L (1996). *Guía para la identificación de los paseriformes europeos*. Sociedad Española de Ornitología, Madrid.
20. KAISER A (1993). A new multicategory classification of subcutaneous fat deposits of songbirds. *Journal of Field Ornithology*, 64:246-255.
21. PINILLA J (2000). *Manual para el anillamiento científico de aves*. SEO/BirdLife y DGCN-MIMAM, Madrid.
22. GOSLER AG, GREENWOOD JJD, BAKER JK & DAVIDSON NC (1998). The field determination of body size and condition in passerines: a report to the British Ringing Committee. *Bird Study*, 45:92-103.
23. LOCKWOOD R, SWADDLE JP & RAYNER JMV (1998). Avian wingtip shape reconsidered: wingtip shape indices and morphological adaptations to migration. *Journal of Avian Biology*, 29:273-292.
24. LIGON JD (1999). *The evolution of avian breeding systems*. Oxford University Press, Oxford.
25. MITCHELL JS & ROBERTSON RJ (1996). Extra nests site occupancy by tree swallows: do floaters avoid nest sites near settled pairs? *Wilson Bulletin*, 108:797-802.

*Received: July 8, 2009*

*Accepted: February 8, 2010*

*Branch editor: Schön Isa*